**Emotion Review** 

Emotion Review

# Bodily contributions to emotion: Schachter's legacy for a psychological constructionist view on emotion

Journal:	Emotion Review
Manuscript ID	EMR-15-845.R2
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	MacCormack, Jennifer; University of North Carolina - Chapel Hill, Psychology and Neuroscience Lindquist, Kristen; University of North Carolina - Chapel Hill, Psychology and Neuroscience
Area/Discipline:	social < psychology
Keywords:	embodiment, core affect, emotion experience, psychological constructionism
Abstract:	Although early emotion theorists posited that bodily changes contribute to emotion, the primary view in affective science over the last century has been that emotions produce bodily changes. Recent findings from physiology, neuroscience, and neuropsychology support the early intuition that body representations can help constitute emotion. These findings are consistent with the modern psychological constructionist hypothesis that emotions emerge when representations of bodily changes are conceptualized as an instance of emotion. We begin by introducing the psychological constructionist approach to emotion. With Schachter as inspiration, we next examine how embodied representations contribute to affective states, and ultimately emotion, with inflammation as a key example. We close by looking forward to future research on how body representations contribute to human experience.



Running head: BODILY CONTRIBUTIONS TO EMOTION
Bodily contributions to emotion: Schachter's legacy for a psychological constructionist view on emotion
Jennifer K. MacCormack & Kristen A. Lindquist
University of North Carolina at Chapel Hill
Address correspondence to:
Kristen Lindquist Department of Psychology and Neuroscience University of North Carolina 321 Davie Hall Chapel Hill, NC 27599 kristen.lindquist@unc.edu

# Abstract

Although early emotion theorists posited that bodily changes contribute to emotion, the primary view in affective science over the last century has been that emotions produce bodily changes. Recent findings from physiology, neuroscience, and neuropsychology support the early intuition that body representations can help constitute emotion. These findings are consistent with the modern psychological constructionist hypothesis that emotions emerge when representations of bodily changes are conceptualized as an instance of emotion. We begin by introducing the psychological constructionist approach to emotion. With Schachter as inspiration, we next examine how embodied representations contribute to affective states, and ultimately emotion, with inflammation as a key example. We close by looking forward to future research on how body representations contribute to human experience.

Keywords: body states, homeostasis, core affect, embodiment, emotion experience,

psychological construction

#### **Emotion Review**

Early theorists posited that body states contribute to emotion experience (e.g., James, 1890). However, the prevailing view in affective science over the last century was that representations of bodily changes are outcomes rather than constituents of emotions. In this view, an emotion, once triggered—either by dedicated neural circuitry (Cannon, 1921; 1927; Ekman & Cordaro, 2011; Panksepp, 1998; Vytal & Hamman, 2010) or a cognitive system (Arnold, 1960; Lazarus, 1966; Roseman, 2011), produces changes in facial and peripheral musculature, as well as patterned changes in the cardiac, respiratory, and neurochemical systems. However, growing evidence suggests that representations of bodily changes may also help *constitute* emotion, by generating the affective states that contribute to emotion experience in the first place (for discussions, see Barrett & Lindquist, 2008; Craig, 2009; Critchley & Nagai, 2012; Damasio, 2001; Laird & Lacasse, 2014; Lindquist, 2013; Oosterwijk et al., 2012).

The idea that representations of body states can constitute, rather than result from, emotions is consistent with a psychological constructionist approach to emotion. We begin our paper by introducing the psychological constructionist approach to emotion and outlining predictions for the role of body representations in emotion. We argue that Stanley Schachter's research played an important, yet frequently unrecognized, role in the legacy of this theory. In particular, Schachter's focus on the constitutive role of body representations in emotion was especially important to the psychological constructionist theory. Building on Schachter's legacy, we next review growing evidence that representations of body states play a constitutive role in emotion and close by examining psychological constructionist predictions for how putatively "non-emotional" bodily changes such as inflammation can contribute to emotion.

A psychological constructionist view: Body representations help constitute emotion

#### **Emotion Review**

Bodily contributions to emotions 4

The idea that body representations help constitute emotion is consistent with modern psychological constructionist views (e.g., Barrett, 2006; Barrett & Russell, 2015; Cunningham, Dunfield, & Stillman, 2013; Lindquist, 2013; Storbeck & Clore, 2008). According to psychological constructionism, an emotion is a highly situated subjective state that emerges from the combination of more basic constitutive parts (for discussions of this approach's causal and measurement models, see Barrett, 2006; 2011).<sup>1</sup> Many constructionist views agree that representations of the body's state serve as one such constitutive part in emotion (e.g., Barrett & Russell, 2015; Cunningham et al., 2013; Lindquist, 2013; Storbeck & Clore, 2008). According to our particular constructionist view, the Conceptual Act Theory (CAT; cf., Barrett, 2006; Barrett et al., 2015; Lindquist, 2013), emotions emerge from causal indicators such as representations of the body's internal state and exteroceptive representations from the current context, as well as attention and conceptual knowledge about emotion categories acquired via prior experiences (for fuller discussion of emotions as emergent variables: Barrett, 2006; 2011; Coan, 2010; Coan & Gonzalez, 2015; Clore & Ortony, 2008). See Figure 1. These elements combine in a given context when representations of bodily changes and exteroceptive sensations are automatically categorized as an instance of an emotion category (e.g., anger vs. fear) using conceptual knowledge about emotions (Barrett, 2015; Lindquist, 2013). Our own approach calls the process by which body representations are made meaningful *situated conceptualization* (cf., Barrett & Russell, 2015; Barrett, Wilson-Mendenhall, & Barsalou, 2015; Lindquist, 2013). Emotions are thus at once bodily, conceptual, and highly situated phenomena (cf., Lindquist & Barrett, 2008; Wilson-Mendenhall et al., 2011).

# Body representations contribute to emotions via core affect

<sup>&</sup>lt;sup>1</sup> Unlike views that assume that emotions are effect indicators (see discussions in Barrett, 2006; Coan, 2010), psychological constructionist views operationalize "emotion" and "emotion experience" as identical, not separate, constructs.

#### **Emotion Review**

All psychological constructionist views agree that *core affect* (sometimes "affect") is a basic, constitutive component of emotions that can be experienced as having some degree of valence and arousal (Barrett, 2006; Clore & Ortony, 2008; Cunningham et al., 2013; Russell, 2003; Russell & Barrett, 1999). Some constructionist views further hypothesize that core affect is derived from representations of changes in the organism's viscera and homeostatic state (Barrett & Bliss-Moreau, 2009; Lindquist, 2013; Russell, 2003; Storbeck & Clore, 2008)<sup>2</sup>. The CAT particularly focuses on how body representations help constitute core affect, and ultimately emotions. Core affect is thought to serve as a homeostatic barometer by bringing disparate representations of bodily information (from the autonomic, proprioceptive, kinesthetic, somatovisceral, and neurochemical systems) together to generate evaluative representations of hedonic valence and arousal (Barrett, 2006; Lindquist, 2013; Lindquist et al., 2012; Russell, 2003).

Critically, insofar as embodied information suffuses and informs every aspect of conscious experience (Barrett & Simmons, 2015; Seth, 2015; Seth, Suzuki, & Critchley, 2012), core affect provides a constantly updating "snapshot" of the body's internal conditions. Core affect may move to the forefront of awareness when there is a noticeable shift (e.g., increases in feelings of arousal, feelings of pleasantness, etc.) or when attention is directed towards our internal state (e.g., when someone asks "how are you?"). The "ambiguous arousal" proposed by Schachter and Singer (1962) is a historical predecessor of the modern-day conception of "core affect".

Core affect is represented by the central nervous system when afferent information from the peripheral nervous system is projected to limbic and paralimbic structures (Craig, 2009;

<sup>&</sup>lt;sup>2</sup> Other views are less explicit about whether core affect derives from body representations (Cunningham et al., 2013; Clore & Ortony, 2013) and focus more on its role as an evaluative state.

#### **Emotion Review**

Bodily contributions to emotions 6

Nauta, 1971). We hypothesize that core affect is sometimes the product of relatively direct representations of body states—for instance, a feeling of arousal associated with perceptions of increased heart rate. However, core affect is perhaps more often a representation of distal bodily causes that may not be directly perceived as specific bodily changes, such as inflammation or low blood sugar, but which contribute to a gestalt representation (e.g., high arousal, unpleasantness, or both).

# Schachter's psychological constructionist legacy

In our view, Schachter's model of emotion (Schachter & Singer, 1962) is a classic example of a psychological constructionist account. However, Schachter's approach was historically aligned with appraisal theories of emotion because the model discussed the role of "meaning making" in emotion. Until recently, it was assumed that any model that invoked "meaning making" in the generation of emotion was necessarily an appraisal model, although this is incorrect (for history, see Gendron & Barrett, 2009). Both appraisal and psychological constructionist approaches focus on meaning making in emotion (see Lindquist, 2013 for a distinction between types of appraisal views). However, what ultimately divides appraisal from psychological constructionist views is that the latter focuses on the role of other fundamental components such as core affect in the generation of emotions. According to constructionist views, core affect is transformed into an emotion experience during situated conceptualization when it is linked to the context and made meaningful as a specific emotion using conceptual knowledge about emotion categories (see Kirkland & Cunningham, 2011; Lindquist, 2013; Wilson-Mendenhall et al., 2011). We believe that Schachter's emphasis on the interactive role between ambiguous physiological arousal from the viscera and the situated conceptualization of that

#### **Emotion Review**

arousal as an emotion during social affiliation thus aligns his approach with modern psychological constructionist accounts.

Schachter's legacy for the psychological constructionist approach stems not only from his focus on the interaction of arousal and meaning making, but also from his particular focus on the bodily representation of arousal. For instance, in his classic study with Singer, he induced arousal via injections of epinephrine, a neurochemical that affects heart rate, respiration, the vasculature, and muscle contraction (Cryer, 1993). Participants' felt arousal in Schachter and Singer's experiment is therefore derived from representations of bodily changes. Even Schachter's lesser-known research had a (at least implicit) focus on the role of body representations in other mental states. For instance, he investigated how body sensations were conceptualized as pain (Nisbett & Schachter, 1966), how representations of unpleasant core affect (i.e., stress) contributed to experiences of hunger (1978, 1982), and how individuals differentially sensed and made meaning of sensations such as hunger (1968, 1971, 1974; with Rodin, 1974). The idea that representations of bodily changes could also be conceptualized as other mental states depending on the context and a person's available conceptual knowledge is consistent with our own psychological constructionist predictions (see Barrett, 2009; Lindquist, 2013; Oosterwijik et al., 2012). For example, core affect derived from body representations could be conceptualized as an emotion (e.g., "anxiety"), as a drive state (e.g., "hunger"), as a somatic complaint (e.g., "an upset stomach"), or as an attitude (e.g., "intense dislike" of something), depending on the situated conceptualization that is made.

Although the specific hypotheses of Schachter and Singer's model were not all supported by early evidence (see Reisenzein, 1983 for review), new evidence substantiates the more general psychological constructionist hypothesis that a person experiences an emotion when core

#### **Emotion Review**

Bodily contributions to emotions 8

affect is made meaningful during situated conceptualization. In particular, accumulating evidence suggests that both peripheral nervous system feedback from the body and central representations of that bodily feedback in the brain are made meaningful as emotion experiences (see Critchley & Harrison, 2013; Harrison et al., 2010 for reviews).

# Body representations contribute to emotion

Although early emotion theorists hypothesized that the body was important for emotion (Duffy, 1957; James, 1884; Sully, 1892), only now with improved methodologies in psychophysiology and neuroscience have researchers been able to test the embodied mechanisms underlying emotion. For example, meta-analyses of neuroimaging studies reveal that a host of brain regions responsible for representing body sensations are active during emotion experience (Kober et al., 2008; Lindquist et al., 2012). One such region is the insula. Neuroanatomical findings suggest that the posterior and mid-insula receive projections from spinal cord lamina representing sensations from the peripheral nervous system and then project this information forward to the anterior insula, a region crucial for integrating specific body representations into a more holistic sense of the body's condition (see Barrett & Simmons, 2015; Craig, 2009; 2011; Critchley, 2004; 2009).

Insula activation, and in particular, anterior insula activation, has been linked to greater *interoceptive sensitivity*—that is, the ability to detect inner bodily changes, such as identifying one's heartbeats accurately (Critchley et al., 2004; Pollatos et al., 2007). Insula activity has also been linked to the experience of other body sensations, such as orgasm (Ortigue et al., 2007), the sensations caused by gastric distention (Vandenbergh et al., 2007; Wang et al., 2008) and pain (Ostrowsky et al., 2002), as well as the perception of other embodied states such as hunger (Del Parigi et al., 2002), breathlessness (Banzett et al., 2000), and craving (Gray & Critchley, 2007).

#### **Emotion Review**

The insula may thus be the hub via which peripheral feedback on the body's current state is translated into core affective representations (Craig, 2009). For instance, individuals with greater interoceptive sensitivity demonstrate greater activity within the insula during emotion experience (Critchley et al., 2004; Gray et al., 2012; Zaki et al., 2012). Similarly, individuals who describe their experiences more in terms of body-based affective terms (e.g., "cry") vs. reflective cognitive terms (e.g., "think") show greater activity within the mid- and anterior insula and primary (SI) and secondary (SII) somatosensory cortex during emotion experience (Saxbe et al., 2013).

The insula, along with brain regions that are directly involved in visceromotor control (e.g., basal ganglia nuclei, periacqueductal gray), project to the orbitofrontal cortex (OFC). Growing evidence suggests that the OFC also contributes to affective experiences (Barrett & Bar, 2009; Bechara et al., 2000; Cunningham et al., 2013; Kringelbach & Rolls, 2004; Lindquist et al., in press; Nauta, 1971; Roy et al., 2012; Schnider et al., 2005) and may help compute the evaluative meaning of body states (Barrett & Simmons, 2015; Cunningham et al., 2013; Lindquist et al., 2013; Lindquist et al., in press).

There are many correlational and experimental studies suggesting that shifts in peripheral physiology alter emotion experience. For example, experimentally manipulating perceived feedback from the body (such as providing false cardiac feedback) can shape how arousing individuals experience their emotions to be (Gray et al., 2012; Gray et al., 2007; Valins, 1966; see Critchley & Garfinkel, 2015 for a review). In correlational studies, individual differences in interoceptive sensitivity are associated with greater reports of high arousal emotions in daily life (e.g., greater anger and excitement vs. sadness and contentment; Barrett et al., 2004). Greater interoceptive sensitivity is also associated with more intense peripheral physiological, self-

#### **Emotion Review**

Bodily contributions to emotions 10

reported, and neural reactions to emotionally evocative images (Herbert et al., 2010; Herbert et al., 2007). For instance, an index of affective intensity called the P300 event-related potential is greater in individuals with greater interoceptive sensitivity (Herbert et al., 2007).

Even intra-individual differences in emotion experience and perceptions appear influenced by body representations. For example, participants experience electric shocks as less aversive when they are delivered prior to ventricular contraction compared to other points within the cardiac cycle (Gray et al., 2009). Shocks cannot produce a change in blood pressure when blood has not yet entered the vasculature; thus, the lack of afferent feedback from the baroreceptors during shock causes emotion experiences to be less intense. The opposite effect is observed when fearful faces are presented supraliminally during the ventricular contraction of the heart—in these cases, fearful faces are more likely to be subjectively seen by participants and more likely to prompt amygdala activity as compared to when they are presented prior to ventricular contraction (Garfinkel et al., 2014). By contrast, activation of the parasympathetic nervous system reduces the affective potency of perceived fearful faces. On trials when the parasympathetic nervous system is activated via pressure to the carotid baroreceptors in the neck, participants view fearful faces as less aversive and show less insula, amygdala, and hippocampus activity when viewing fearful faces (Makovac et al., 2015). Activation of the parasympathetic nervous system presumably induces experiences of calm and inhibits activation of highly arousing feelings produced by the sympathetic nervous system.

Conditions in which internal body representations are impaired or attenuated more specifically point to the body's role in emotion experience. Pure autonomic failure (PAF) is a neurodegenerative disease linked to autonomic denervation, resulting in decreased transmission of afferent information from the peripheral nervous system to the brain (Critchley, Mathias, &

#### **Emotion Review**

Dolan, 2002). PAF is also associated with decreased peripheral physiological activity, such as decreases in skin conductance (Magnifico et al., 1998), heart rate (Mathias & Bannister, 1999), and catecholamine levels (e.g., epinephrine, norepinephrine, and dopamine; Meredith et al., 1991). Critically, with reductions in sympathetic changes and autonomic feedback, individuals with PAF report experiencing less stress when performing a difficult math task in front of peers than do patient controls (Critchley, Mathias, & Dolan, 2001). These same individuals also demonstrate decreased insula activity and possess less awareness about the aversive nature of stimuli during classical conditioning (Critchley et al., 2002). With significantly reduced afferent information from the body, individuals seem less able to acquire the representation that stimuli are aversive.

Even healthy aging causes some peripheral attenuation (Levenson et al., 1991; Neiss et al., 2009; Tsai, Levenson, & Carstensen, 2000) and is thought to produce emotion experiences that are less linked to representations of the body's current state and more associated with external, situational cues (cf., Mendes, 2010). These findings may in part account for the age-related changes observed in emotion such as older individuals' increased facility in emotion regulation (Gross et al., 1997; Urry & Gross, 2010). It is perhaps relatively easier to regulate one's emotions when you don't experience intense bodily changes.

The fact that the elderly and patients with PAF still retain some emotional abilities even in the presence of changes in body representation is itself ultimately consistent with our psychological constructionist account. Body representations are not the only constitutive element in emotion; thus, maintained emotional abilities may stem from the maintenance of other elements. First, in the cases of PAF and aging, both proprioception and some somatovisceral representations appear to remain intact, even as peripheral representations decline (Critchley et al., 2002; Mendes, 2010). Second, and most critically, patients with PAF and individuals undergoing healthy aging still possess the conceptual knowledge about emotion experiences that they have amassed over their lifetimes. Even in the absence of robust online representations of bodily changes, centrally represented conceptual knowledge about the internal and external cues associated with emotions could potentially be used to compensate for peripheral declines, so that individuals still retain somewhat normal emotion experiences. Consistent with this hypothesis, patients with PAF are still able to engage in normative emotion recognition, motivational decision-making, theory of mind, and social cognition (Heims et al., 2004). Similarly, although recent evidence suggests that patients with high spinal cord injuries may experience changes in the intensity or specificity of emotions following their injuries (Nicotra et al., 2006; Pistoia et al., 2015), patients retain accurate conceptual knowledge about emotions after their injuries (Deady et al., 2010).

Unlike neo-Jamesian interpretations of the body's role in emotion (for discussion, see Reisenzein & Stephan, 2014), we are not arguing that body representations are alone sufficient for emotion. Current body representations may not even be strictly necessary after a person has acquired years of conceptual knowledge that can be simulated by the brain during emotion experience (after all, spinal cord injury patients can simulate the experience of limb pain and pleasure in the absence of real sensations; Sjogren & Egberg, 1983). Of course, the purest test of the body's constitutive role in emotion would be to observe individuals who had never experienced the representation of bodily changes from birth, but finding such cases are difficult, if not impossible in humans.

Whereas much research has examined the peripheral physiological and visceral contributions to emotion, far less research has focused on how other bodily systems contribute to

http://mc.manuscriptcentral.com/emotion-review

#### **Emotion Review**

emotion experience. We turn to this research now. A unique prediction of the psychological constructionist view is that myriad homeostatic bodily processes—such as immune responses that induce inflammation, hormonal fluctuations, and even broader "drives" such as hunger, fatigue, and sexual arousal—feed into core affective representations that then may become conceptualized as instances of discrete emotions. If conceptualized as an instance of emotion, these bodily changes could contribute to emotions in ways that the field has little considered, breaking down the traditional distinction between "motivational states" vs. "emotions," and even "the immune system" vs. "the affective system." As a case in point, we explore preliminary evidence suggesting that inflammation is one source of bodily change that can shift individuals' core affective states, and then hypothesize how such inflammation-induced shifts in core affect might be experienced as specific discrete emotions.

# Inflammation contributes to core affect and emotion

Although the "immune system" and the "affective system" have been traditionally viewed as distinct, growing evidence suggests that inflammatory responses related to immune function also cause shifts in core affect, and that some of the same neurochemicals involved in immune responses can generate affective changes. Herein, we discuss the inflammatory response in particular, and its relation to and potential impact on emotion.

# Neurochemical mechanisms of inflammation

Inflammation is an immune response to harmful stimuli that occurs across a wide range of disorders (see Iwata, Ota, & Duman, 2013) as well as in response to tissue trauma, chemotherapy, surgery, and social stressors (Brydon et al., 2004; Cole et al., 2007; 2012; O'Conner et al., 2014). Both local and systemic inflammation involve pro-inflammatory cytokines, which are associated with a variety of physiological changes and somatic sensations; we thus suggest that pro-inflammatory cytokines may serve as one mechanism by which inflammation feeds into core affect, and ultimately, emotion.

# Inflammatory contributions to core affect

Research increasingly demonstrates that the presence of pro-inflammatory cytokines correlates with and induces the experience of heightened and persistent unpleasant core affect. Regular administration of cytokines in humans is correlated with increased reports of general unpleasantness (Raison et al., 2010; Reichenberg et al., 2001; Schiepers et al., 2005) and injections of bacteria that promote an inflammatory response in the immune system increase negative mood (Brydon et al., 2008; Strike et al., 2004; Wright et al., 2005). Induced inflammation can also produce high arousal states (e.g., irritability, insomnia, hyperactivity) that can be experienced as either positive or negative affect (Constant et al., 2005). In contrast, acetaminophen, a drug that may dampen central nervous system inflammation, reduces the selfreported intensity of positive and negative states (Durso, Lutrell, & Way, 2015).

Inflammation's impact on core affect is further suggested by neuroimaging studies that examine central nervous system activity in the presence of pro-inflammatory cytokines (for full review, see Dantzer et al., 2008). For instance, typhoid- (vs. placebo-) injected participants experience increased activation in the neural regions supporting the generation and representation of core affect including the brainstem, thalamus, amygdala, anterior cingulate, and the anterior insula (Harrison et al., 2009a, b). Critically, changes in core body temperature or self-reported somatic symptoms were not experienced, suggesting that shifts in negative core affect do not merely result from feelings of sickness.

## Inflammatory contributions to discrete emotion

#### **Emotion Review**

The key hypothesis that we put forward, inspired by psychological constructionism and ultimately Schachter's work, is that when an inflammatory response leads to core affective changes, it is then possible for those inflammatory-induced representations to become conceptualized as a discrete emotion. The dynamics of when and how these core affective shifts are made meaningful as specific emotions likely depends as much on the context and a person's available conceptual knowledge as the core affective shifts themselves (e.g., unpleasantness might be experienced as fatigue in one context, but sadness in another).

In the case of chronic inflammation, individuals likely experience persistent shifts in core affect that track the production cycles of pro-inflammatory cytokines (such as in depression, which has long been linked with chronic, systemic inflammation; Dowlati et al., 2010; Howren et al., 2009). Shifts in pro-inflammatory cytokines might be conceptualized as specific emotions (e.g., sadness) when the context makes knowledge about sadness more accessible (e.g., in the face of a failed job performance), but not when sadness is relatively less accessible (e.g., when watching an enjoyable movie with a friend). Chronic inflammation might lead to the chronic accessibility of negative emotion concepts which could perpetuate the cycle of negative affectivity in some cases of chronic inflammation, thus increasing the likelihood of depression (especially in combination with adverse life events). By contrast, acute inflammation might cause only a temporary fluctuation in core affect, which may make individuals access situationspecific emotion concepts in that instance and then experience only a relatively short-lived experience of emotion. Either acute or chronic shifts in core affect may interact with the context and conceptual knowledge, essentially "turning up the dial" on core affect and subsequent emotion experiences that might have been likely to occur in that context anyway. For instance,

experiencing a loss while experiencing an acute shift in core affect due to the flu might increase the intensity of sadness experienced.

No research to date has explicitly assessed how making meaning of one's core affective state, as induced by inflammation, can result in the experience of different discrete emotions. However, some evidence is suggestive. For instance, in the United States, people who have greater chronic inflammation are also chronically more likely to express anger; in Japan, the inverse is the case (Kitayama et al., 2015). One interpretation is that individuals who have greater inflammation (perhaps due to lifestyle) are more likely to conceptualize their feelings of inflammation as anger in U.S. contexts than in Japanese contexts. Since the U.S. concept of anger emphasizes anger as an emotion that should be expressed when feeling strong unpleasant, high arousal affect, U.S. participants may be more likely to conceptualize their inflammationrelated unpleasantness and high arousal as feelings of anger. By contrast, the Japanese concept of anger condemns expressions of anger (Boiger et al., 2013) and the Japanese make interpersonal adjustments to reduce the likelihood of experiencing anger during conflict (Trommsdorff & Kornadt, 2003). Thus, in Japan, greater inflammation may be less readily associated with highly unpleasant, activated feelings; feelings associated with inflammation may in turn be less likely to become conceptualized as an instance of anger.

In other contexts, individuals experiencing negativity and tension due to inflammation might be more likely to conceptualize their state as an instance of anxiety (see Moons & Shields, 2015). For instance, individuals given inflammation-inducing endotoxin vs. placebo show heightened amygdala response to threatening, but not non-threatening, faces and images (Inagaki et al., 2012). Amygdala activity is associated with uncertainty (Whalen, 2007) and individuals with anxiety disorders have heightened amygdala activity (Etkin & Wager, 2007). As in previous

#### **Emotion Review**

studies, inflammation was constructed as an experience of anxiety when participants were faced with threatening faces and scenes, above and beyond any feelings of sickness that participants reported (Inagaki et al., 2012).

The findings reviewed above are preliminary, and the studies themselves were not designed to answer questions about how individuals are conceptualizing inflammation-related core affect as instances of discrete emotion across different contexts. However, these findings provide promising evidence in line with a psychological constructionist approach that body representations can "feed into" core affect and discrete emotion experiences. Thinking of immune responses as only one of several bodily sources that contribute to shifts in core affect may ultimately be the most parsimonious way for framing the immune system's relation to the affective system.

### Conclusion

The evidence on inflammation we reviewed suggests that representations of body states, including more basic homeostatic processes not typically associated with emotion, feed into core affect and can become conceptualized as emotions. We can thank Schachter for inspiring a psychological constructionist approach that emphasizes the body's constitutive role in emotion. Extending the body's role in emotion to include inter- and intra-individual variation in homeostatic processes could, if further explored, deepen our understanding of why emotion experiences vary so much within and between individuals, and will further clarify what we know about the link between emotions, health, and wellbeing.

We highlighted research suggesting that inflammatory responses drive affective changes that might, in conjunction with a person's conceptual knowledge, be conceptualized as specific discrete emotions. These findings, if further tested, will clarify how putatively "non-emotional"

#### **Emotion Review**

Bodily contributions to emotions

body states (e.g., immune responses, hunger, fatigue) could shape emotion, and more specifically, how longstanding changes in representations of body-based information due to illness, trauma, or even normative development (e.g., puberty, aging) impact individuals' core affective states and how they conceptualize core affect as experiences of discrete emotions. For example, patients with inflammation-inducing disorders may be at greater risk for developing emotion-related psychopathologies (e.g., depression, anxiety, more labile anger) that may feed back to negatively shape physical health outcomes.

Although we did not review evidence beyond inflammation, we suggest that other homeostatic body states will also contribute to shifts in emotion experience. For instance, hungry individuals appear more likely to experience anger. It is typically assumed that hunger depletes individuals' regulatory resources (cf., Bushman et al., 2014), but it is also possible that hunger helps create anger when individuals conceptualize the unpleasant, high arousal changes induced by hunger as an instance of anger if given a frustrating situation. Similarly, fatigue could be conceptualized as instances of loneliness, depression, or sadness. Future research should investigate these questions and assess how homeostatic bodily changes shape a person's core affect and ultimately emotion. Such work would clarify how bodily changes help constitute and create variation in emotion experience. Thanks in part to Schachter's legacy, the field of affective science is further exploring these questions about how the body constitutes emotions and the mind more generally.

#### **Emotion Review**

# References

- Arnold, M.B. (1960). Emotion and personality. New York, NY: Columbia University Press.
- Banzett, R.B., Mulnier, H.E., Murphy, K., Rosen, S.D., Wise, R.J., & Adams, L. (2000). Breathlessness in humans activates insular cortex. *Neuroreport*, *11*, 2117-2120.
- Barrett, L.F. (2006). Solving the emotion paradox: Categorization and the experience of emotion. *Personality and Social Psychology Review, 10*, 20-46.
- Barrett, L.F. (2009). The future of psychology: Connecting mind to brain. *Perspectives on Psychological Science*, *4*, 326–339.
- Barrett, L.F (2011). Bridging token identity theory and supervenience theory through psychological construction. *Psychological Inquiry, 22*, 115-127.
- Barrett, L.F. (2015). Ten common misconceptions about the psychological construction of emotion. Chapter in L. F. Barrett and J. A. Russell (Eds.) *The psychological construction of emotion* (p. 45-79). New York: Guilford.
- Barrett, L.F. & Simmons, W.K. (2015). Interoceptive predictions in the brain. *Nature Reviews Neuroscience*, *16*, 419-429.
- Barrett, L.F., & Bar, M. (2009). See it with feeling: Affective predictions in the human brain. *Royal Society Phil Trans B, 364,* 1325-1334.
- Barrett, L.F., & Bliss-Moreau, E. (2009). Affect as a psychological primitive. Advances in Experimental Social Psychology, 41, 167-218.
- Barrett, L.F., & Lindquist, K. (2008). The embodiment of emotion. In G. Semin & E. Smith (Eds.), *Embodied grounding: Social, cognitive, affective, and neuroscience approaches* (pp. 237-262). New York: Cambridge University Press.
- Barrett, L.F., & Russell, J.A. (2015). An introduction to psychological construction. In L. F. Barrett and J. A. Russell (Eds.), *The psychological construction of emotion* (pp. 1-17).

New York: Guilford.

- Barrett, L.F., Quigley, K., Bliss-Moreau, E., & Aronson, K.R. (2004). Interoceptive sensitivity and reports of emotional experience. *Journal of Personality and Social Psychology*, 87, 684-697.
- Barrett, L.F., Wilson-Mendenhall, C.D., & Barsalou, L.W. (2015). The conceptual act theory: a road map. Chapter in L.F. Barrett and J.A. Russell (Eds.), *The psychological construction of emotion* (pp. 83-110). New York: Guilford.
- Bechara, A., Damasio, H., & Damasio, A.R. (2000). Emotion, decision-making and the orbitofrontal cortex. *Cerebral Cortex*, 10, 295–307.
- Boiger, M., Mesquita, B., Uchida, Y., & Barrett, L.F. (2013). Condoned or condemned: The situational affordance of anger and shame in the United States and Japan. *Personality and Social Psychology Bulletin, 39*, 540-553.
- Brydon, L., Edwards, S., Mohamed-Ali, V., & Steptoe, A. (2004). Socioeconomic status and stress-induced increases in interleukin-6. *Brain, Behavior, & Immunity, 18,* 281–290.
- Brydon, L., Harrison, N.A., Walker, C., Steptoe, A., & Critchley, H.D. (2008). Peripheral inflammation is associated with altered substantia nigra activity and psychomotor slowing in humans. *Biological Psychiatry*, *63*, 1022–1029.
- Bushman, B.J., DeWall, C.N., Pond Jr., R.S., & Hanus, M.D. (2014). Low glucose relates to greater aggression in married couples. *Proceedings of the National Academy of Sciences*, U.S.A., 111, 6254-6257.
- Cannon, W.B. (1921). The James–Lange theory of emotions: A critical examination and an alternative theory. *American Journal of Psychology*, *38*, 106–124.

Cannon W.B. (1927). Bodily changes in pain, hunger, fear and rage: An account of recent

#### **Emotion Review**

researches into the function of emotional excitement. New York: D. Appleton.

- Clore, G.L., & Ortony, A. (2008). Appraisal theories: How cognition shapes affect into emotion.
  In J. M. Haviland-Jones, M. Lewis, & L. F. Barrett (Eds.), *Handbook of emotions* (Vol. 3, pp. 628–642). New York, NY: Guilford.
- Clore, G.L., & Ortony, A. (2013). Psychological construction in the OCC model of emotion. *Emotion Review, 5*, 335-343.

Coan, J.A. (2010). Emergent ghosts of the emotion machine. *Emotion Review*, 2, 274-285.

- Coan, J.A. & Gonzalez, M.Z. (2015). Emotions as emergent variables. In L.F. Barrett & J.A.
   Russell (Eds.), *The Psychological Construction of Emotion* (pp. 209-226). New York: The Guilford Press.
- Cole, S.W., Conti, G., Arevalo, J.M.G., Ruggiero, A.M., Heckman, J.J., & Suomi, S.J. (2012)
   Transcriptional modulation of the developing immune system by early life social
   adversity. *Proceedings from the National Academy of Sciences USA 109*, 20578–20583.
- Cole, S.W., Hawkley, L.C., Arevalo, J.M.G., Sung, C.Y., Rose, R.M., & Cacioppo, J.T. (2007). Social regulation of gene expression in human leukocytes. *Genome Biology*, *8*, R189.
- Constant, A., Castera, L., Dantzer, R., Couzigou, P., de Ledinghen, V., Demotes-Mainard, J., & Henry, C. (2005). Mood alterations during interferon-alpha therapy in patients with chronic hepatitis C: Evidence for an overlap between manic/hypomanic and depressive symptoms. *Journal of Clinical Psychiatry*, *66*, 1050-1057.
- Craig, A.D. (2009). How do you feel—now? The anterior insula and human awareness. *Nature Reviews Neuroscience*, *10*, 59-70.
- Craig, A.D. (2011). Significance of the insula for the evolution of human awareness of feelings from the body. *Annals of the New York Academy of Sciences, 1225*, 72-82.

- Critchley, H.D. (2004). The human cortex responds to an interoceptive challenge. *Proceedings* from the National Academy of Sciences, 101, 6333–6334.
- Critchley, H.D. (2009). Psychophysiology of neural, cognitive, and affective integration: fMRI and autonomic indicants. *International Journal of Psychophysiology*, *73*, 88-94.
- Critchley, H.D., & Garfinkel, S.N. (2015). Interactions between visceral afferent signaling and stimulus processing. *Frontiers in Neuroscience*, *9*, 286.
- Critchley, H.D., & Harrison, N.A. (2013). Visceral influences on brain and behavior. *Neuron*, 77, 624-638.
- Critchley, H.D., & Nagai, Y. (2012). How emotions are shaped by bodily states. *Emotion Review*, *4*, 163-168.
- Critchley, H.D., Mathias, C.J., & Dolan, R.J. (2001). Neuroanatomical basis for first- and second-order representations of bodily states. *Nature Neuroscience*, *4*, 207-212.
- Critchley, H.D., Mathias, C.J., & Dolan, R.J. (2002). Fear conditioning in humans: The influence of awareness and autonomic arousal on functional neuroanatomy. *Neuron, 33*, 653-663.
- Critchley, H.D., Wiens, S., Rotshtein, P., Ohman, A., & Dolan, R.J. (2004). Neural systems supporting interoceptive awareness. *Nature Neuroscience*, *7*, 189–95.
- Cryer, P.E. (1993). Adrenaline: A physiological metabolic regulatory hormone in humans. *International Journal of Obesity and Related Metabolic Disorders, 17*, S43-46.
- Cunningham, W.A., Dunfeld, K.A., & Stillman, P. (2013). Emotional states from affective dynamics. *Emotion Review*, *5*, 344-355.
- Damasio, A.R. (2001). Fundamental feelings. Nature, 413, 781.
- Dantzer, R., O'Connor, J.C., Freund, G.G., Johnson, R.W., Kelley, K.W. (2008). From inflammation to sickness and depression: When the immune system subjugates the brain.

Nature Reviews Neuroscience, 9, 46–56.

- Deady, D.K., North, N.T., Allan, D., Law Smith, M.J., & O'Carroll, R.E. (2010). Examining the effect of spinal cord injury on emotional awareness, expressivity, and memory for emotional material. *Psychology, Health & Medicine, 15*, 406-419.
- Del Parigi, A., Chen, K., Gautier, J.F., Salbe, A.D., Pratley, R.E., Ravussin, E., Reiman, E.M., & Tataranni, P.A. (2002). Sex differences in the human brain's response to hunger and satiation. *American Journal of Clinical Nutrition*, 75, 1017-1022.
- Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E.K., & Lanctot, K.L.
  (2010). A meta-analysis of cytokines in major depression. *Biological Psychiatry*, 67, 446-457.
- Duffy, E. (1957). The psychological significance of the concept of arousal or activation. *Psychological Review, 64,* 265-275.
- Durso, G.R.O., Luttrell, A., & Way, B. (2015). Over-the-counter relief from pains and pleasures alike: Acetaminophen blunts evaluation sensitivity to both negative and positive emotional stimuli. *Psychological Science*, *26*, 750-758.
- Ekman, P., & Cordaro, D. (2011). What is meant by calling emotions basic. *Emotion Review*, *3*, 364–370.
- Etkin, A., & Wager, T.D. (2007). Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*, 164, 1476-1488.
- Garfinkel, S.N., Minati, L., Gray, M.A., Seth, A.K., Dolan, R.J., & Critchley, H.D. (2014). Fear from the heart: Sensitivity to fear stimuli depends on individual heartbeats. *The Journal of Neuroscience, 34*, 6573-6582.

- Gray, M.A., & Critchley, H.D. (2007). Interoceptive basis of craving. Neuron, 54, 183-186.
- Gray, M.A., Beacher, F., Minati, L., Nagai, Y., Kemp, A.H., Harrison, N., & Critchley, H.D.
  (2012). Emotional appraisal is influenced by cardiac afferent information. *Emotion*, *12*, 180-191.
- Gray, M.A., Harrison, N.A., Wiens, S., & Critchley, H.D. (2007). Modulation of emotional appraisal by false physiological feedback during fMRI. *PLoS ONE, 2,* e546.
- Gray, M.A., Minati, L., Paoletti, G., & Critchley, H.D. (2009). Baroreceptor activation attenuates attentional effects on pain-evoked potentials. *Pain, 151,* 853-861.
- Gross, J.J., Carstensen, L.C., Pasupathi, M., Tsai, J., Gottestam, K., & Hsu, A.Y.C. (1997).
  Emotion and aging: Experience, expression, and control. *Psychology and Aging*, *12*, 590-599
- Harrison, N.A., Brydon, L., Walker, C., Gray, M.A., Steptoe, A., & Critchley, H.D. (2009a).
   Inflammation causes mood changes through alterations in subgenual cingulate activity and mesolimbic connectivity. *Biological Psychiatry*, 66, 407-414.
- Harrison, N.A., Brydon, L., Walker, C., Gray, M.A., Steptoe, A., Dolan, R.J., & Critchley, H.D.
  (2009b). Neural origins of human sickness in interoceptive responses to inflammation. *Biological Psychiatry*, 66, 415-422.
- Harrison, N.A., Gray, M.A., Gianaros, P.J., & Critchley, H.D. (2010). The embodiment of emotional feelings in the brain. *Journal of Neuroscience*, 30, 12878-12884.
- Heims, H.C., Critchley, H.D., Dolan, R., Mathias, C.J., & Cipolotti, L. (2004). Social and motivational functioning is not critically dependent on feedback of autonomic responses: Neuropsychological evidence from patients with pure autonomic failure. *Neuropsychologia*, 42, 1979-1988.

- Herbert, B.M., Pollatos, O., & Schandry, R. (2007). Interoceptive sensitivity and emotion processing: An EEG study. *International Journal of Psychophysiology*, 65, 214-227.
- Herbert, B.M., Pollatos, O., Flor, H., Enck, P., & Schandry, R. (2010). Cardiac awareness and autonomic cardiac reactivity during emotional picture viewing and mental stress. *Psychophysiology*, 47, 342-354.
- Howren, M.B., Lamkin, D.M., Suls, J. (2009). Associations of depression with C-reactive protein, IL-1, and IL-6: A meta-analysis. *Psychosomatic Medicine*, *71*, 171–186.
- Inagaki, T.K., Muscatell, K.A., Irwin, M. R., Cole, S.W., & Eisenberger, N.I. (2012). Inflammation selectively enhances amygdala activity to socially threatening images. *Neuroimage, 59*, 3222-3226.
- Iwata, M., Ota, K.T., & Duman, R.S. (2013). The inflammasome: Pathways linking psychological stress, depression, and systemic illnesses. *Brain, Behavior, and Immunity*, 31, 105-114.
- James, W. (1884). What is an emotion? Mind, 34, 188–205.
- James, W. (1890). The principles of psychology. New York, NY: Holt.
- Kirkland, T., & Cunningham, W.A. (2011). Neural basis of affect and emotion. *Wiley Interdisciplinary Reviews: Cognitive Science, 2,* 656-665.

Kitayama, S., Park, J., Boylan, J.M., Miyamoto, Y., Levine, C.S., Markus, H.R., Karasawa, M.,
Coe, C.L., Kawakami, N., Love, G.D., & Ryff, C.D. (2015). Expression of anger and ill
health in two cultures: An examination of inflammation and cardiovascular risk. *Psychological Science*, 26, 211-220.

Kober, H., Barrett, L.F., Joseph, J., Bliss-Moreau, E., Lindquist, K.A., & Wager, T.D. (2008). Functional networks and cortical-subcortical interactions in emotion: A meta-analysis of neuroimaging studies. Neuroimage, 42, 998-1031.

- Kringelbach, M.L., & Rolls, E.T. (2004). The functional neuroanatomy of the human orbitofrontal cortex: Evidence from neuroimaging and neuropsychology. *Progress in Neurobiology*, 72, 341-372.
- Laird, J.D., & Lacasse, K. (2014). Bodily influences on emotional feelings: Accumulating evidence and extensions of William James' theory of emotion. *Emotion Review, 6,* 27-34.

Lazarus, R.S. (1966). Psychological stress and the coping process. New York: McGraw-Hill.

- Levenson, R.W., Carstensen, L.L., Friesen, W.V., & Ekman, P. (1991). Emotion, physiology, and expression in old age. *Psychology and Aging*, *6*, 28–35.
- Lindquist, K.A. (2013). Emotions emerge from more basic psychological ingredients: A modern psychological constructionist approach. *Emotion Review*, *5*, 356-368.
- Lindquist, K.A., & Barrett, L.F. (2008). Constructing emotion: The experience of fear as a conceptual act. *Psychological Science*, *19*, 898-903.
- Lindquist, K.A., MacCormack, J.K., & Shablack, H. (2015). The role of language in emotion: Predictions from psychological constructionism. *Frontiers in Psychology*, *6*, 444.
- Lindquist, K.A., Satpute, A.B., Wager, T.D., Weber, J., & Barrett, L.F. (in press). The brain basis of positive and negative affect: Evidence from a meta-analysis of the human neuroimaging literature. *Cerebral Cortex*.
- Lindquist, K.A., Wager, T.D., Kober, H., Bliss-Moreau, E., & Barrett, L.F. (2012). The brain basis of emotion: A meta-analytic review. *Behavioral and Brain Sciences*, *35*, 121-143.
- Magnifico, F., Misra, V.P., Murray, N.M., & Mathias, C.J. (1998). The sympathetic skin response in peripheral autonomic failure—evaluation in pure failure pure cholinergic dysautonomia and dopamine-beta-hydroxylase deficiency. *Clinical Autonomic Research,*

 8, 133–138.

# Makovac, E., Garfinkel, S.N., Bassi, A., Basile, B., Macaluso, E., Cercignani, M., Calcagnini, G., Mattei, E., Agulliu, D., Cortelli, P., Caltagirone, C., Bozzali, M., & Critchley, H.D. (2015). Effect of parasympathetic stimulation on brain activity during appraisal of fearful expressions. *Neuropsychopharmacology*, *40*, 1649-1658.

- Mathias, C.J., & Bannister, R. (1999). Investigation of autonomic disorders. Mathias, C. J. &
   Bannister, R. (Eds.), *Autonomic Failure: A Textbook of Clinical Disorders of the Autonomic Nervous System* (4th edition), Oxford University Press, Oxford, pp. 169–195.
- Mendes, W.B. (2010). Weakened links between mind and body in older age: The case for maturational dualism in the experience of emotion. *Emotion Review*, *2*, 1–5.
- Meredith, I.T., Esler, M.D., Cox, H.S., Lambert, G.W.L., Jennings, G.L., & Eisenhofer, G.
  (1991). Biochemical evidence of sympathetic denervation of the heart in pure autonomic failure. *Clinical Autonomic Research*, *1*, 187-194.
- Moons, W.G., & Shields, G.S. (2015). Anxiety, not anger, induces inflammatory activity: An avoidance/approach model of immune system activation. *Emotion*, *15*, 463-476.
- Nauta, W.J.H. (1971). The problem of the frontal lobe: A reinterpretation. *Journal of Psychiatric Research*, *8*, 167-187.
- Neiss, M.B., Leighland, L.A., Carlson, N.E., & Janowsky, J.S. (2009). Age differences in perception and awareness of emotion. *Neurobiology of Aging*, *30*, 1305-1313.
- Nicotra, A., Critchely, H.D., Mathias, C.J., & Dolan, R.J. (2006). Emotional and autonomic consequences of spinal cord injury explored using functional brain imaging. *Brain*, 129, 718-728.

Nisbett, R.E. & Schachter, S. (1966). Cognitive manipulation of pain. Journal of Experimental

Social Psychology, 2, 227-236.

- O'Conner, M.F., Schultze-Florey, C.R., Irwin, M.R., Arevalo, J.M., & Cole, S.W. (2014). Divergent gene expression responses to complicated grief and non-complicated grief. *Brain, Behavior, and Immunity, 37*, 78-83.
- Oosterwijk, S., Lindquist, K.A., Anderson, E., Dautoff, R., Moriguchi, Y., & Barrett, L.F. (2012). States of mind: Emotions, body feelings, and thoughts share distributed neural networks. *Neuroimage*, *62*, 2110-2128.
- Ortigue, S., Grafton, S.T., & Bianchi-Demicheli, F. (2007). Correlation between insula activation and self-reported quality of orgasm in women. *NeuroImage*, *37*, 551-560.
- Ostrowsky, K., Magnin, M., Ryvlin, P., Isnard, J., Geunot, M., Mauguiere, F. (2002). Representation of pain and somatic sensation in the human insula: A study of responses to direct electrical cortical stimulation. *Cerebral Cortex*, *12*, 376-385.
- Panksepp, J. (1998) *Affective neuroscience: The foundations of human and animal emotions.* Oxford University Press.
- Pistoia, F., Carolei, A., Sacco, S., Conson, M., Pistarini, C., Cazzulani, B., Stewart, E.J.,
  Franceschini, M., & Sara, M. (2015). Contributions of interoceptive information to
  emotional processing: Evidence from individuals with spinal cord injury. *Journal of Neurotrauma*, epub ahead of print.
- Pollatos, O., Schandry, R., Auer, D.P., & Kaufmann, C. (2007). Brain structures mediating cardiovascular arousal and interoceptive awareness. *Brain Research*, *1141*, 178-187.
- Raison, C.L., Borisov, A.S., Woolwine, B.J., Massung, B., Vogt, G., & Miller, A.H. (2010).
   Interferon-alpha effects on diurnal hypothalamic-pituitary-adrenal axis activity:
   Relationship with proinflammatory cytokines and behavior. *Molecular Psychiatry*, 15,

535-547.

- Reichenberg, A., Yirmiya, R., Schuld, A., Kraus, T., Haack, M., Morag, A., & Pollmacher, T.
   (2001). Cytokine-associated emotional and cognitive disturbances in humans. *Archives of General Psychiatry*, 58, 445-452.
- Reisenzein, R. (1983). The Schachter theory of emotion: Two decades later. *Psychological Bulletin, 94,* 293-264.
- Reisenzein, R., & Stephan, A. (2014). More on James and the physical basis of emotion. *Emotion Review, 6,* 35-46.
- Roseman, I.J. (2011). Emotional behaviors, emotivational goals, emotion strategies: Multiple levels of organization integrate variable and consistent responses. *Emotion Review*, *3*, 1– 10.
- Roy, M., Shohamy, D., & Wager, T.D. (2012). Ventromedial prefrontal-subcortical systems and the generation of affective meaning. *Trends Cognitive Science*, 16, 147–156.
- Russell, J.A. (2003). Core affect and the psychological construction of emotion. *Psychological Review*, *110*, 145–172.
- Russell, J.A., & Barrett, L.F. (1999). Core affect, prototypical emotional episodes, and other things called emotion: Dissecting the elephant. *Journal of Personality and Social Psychology*, 76, 805–819.
- Saxbe, D.E., Yang, X., Borofsky, L.A., & Immordino-Yang, M.H. (2013). The embodiment of emotion: Language use during the feeling of social emotions predicts cortical somatosensory activity. *Social Cognitive and Affective Neuroscience*, 8, 806-812.

Schachter, S. (1968). Obesity and eating. Science, 161, 751-756.

- Schachter, S. (1971). Some extraordinary facts about obese humans and rats. *American Psychologist, 26,* 129-144.
- Schachter, S. (1974). Appetite regulation in obese subjects. *Hormone and Metabolic Research, 4,* 88-93.
- Schachter, S. (1978). Pharmacological and psychological determinants of cigarette smoking. Annals of Internal Medicine, 88, 104-114.
- Schachter, S. (1982). Recidivism and self-cure of smoking and obesity. *American Psychologist, 37*, 436-444.
- Schachter, S., & Rodin, J. (1974). Obese humans and rats. Hillsdale, NJ: Erlbaum.
- Schachter, S., & Singer, J.E. (1962). Cognitive, social, and physiological determinants of emotional state. *Psychological Review*, *69*, 378-399.
- Schiepers, O.J., Wichers, M.C., Maes, M. (2005). Cytokines and major depression. *Progress in Neuro-psychopharmacology and Biological Psychiatry*, 29, 201–217.
- Schnider, A., Treyer, V., & Buck, A. (2005). The human orbitofrontal cortex monitors outcomes even when no reward is at stake. *Neuropsychologia*, *43*, 316-323.
- Seth, A.K. (2015). The cybernetic Bayesian brain: From interoceptive inference to sensorimotor contingencies. In T. Metzinger & J. M. Windt (Eds.), *Open MIND*: 35(T). Frankfurt am Main: MIND Group.
- Seth, A.K., Suzuki, K., & Critchley, H.D. (2012). An interoceptive predictive coding model of conscious presence. *Frontiers in Psychology*, 2, 395.
- Sjogren, K., & Egberg, K. (1983). The sexual experience in younger males with complete spinal cord injury. *Scandinavian Journal of Rehabilitation Medicine*, *9*, 189-194.

- Storbeck, J., & Clore, G.L. (2008). Affective arousal as information: How affective arousal influences judgments, learning, and memory. *Social and Personality Psychology Compass, 2*, 1824-1843.
- Strike, P.C., Wardle, J., & Steptoe, A. (2004). Mild acute inflammatory stimulation induces transient negative mood. *Journal of Psychosomatic Research*, *57*, 189–194.

Sully, J. (1892). *Outlines of psychology*. New York: Appleton & Company.

- Trommsdorff, G., & Kornadt, H. (2003). Parent-child relations in cross-cultural perspective. In L. Kuczynski (Ed.), *Handbook of dynamics in parent-child relations* (pp. 271-305). Thousand Oaks, CA: SAGE.
- Tsai, J.L., Levenson, R.W., & Carstensen, L.L. (2000). Autonomic, expressive, and subjective responses to emotional films in older and younger Chinese American and European American adults. *Psychology and Aging*, 15, 684-693.
- Urry, H.L. & Gross, J.J. (2010). Emotion regulation in older age. *Current Directions in Psychological Science*, 19, 352-357.
- Valins, S. (1966). Cognitive effects of false heart-rate feedback. *Journal of Personality and Social Psychology, 4,* 400-408.
- Vandenbergh, J., Dupont, P., Fischler, B., Bormans, G., Persoons, P., Janssens, J., & Tack, J. (2005). Regional brain activation during proximal stomach distention in humans: A positron emission tomography study. *Gastroenterology*, *128*, 564-573.
- Vytal, K., & Hamann, S. (2010). Neuroimaging support for discrete neural correlates of basic emotions: A voxel-based meta-analysis. *Journal of Cognitive Neuroscience, 22*, 2864-2885.

Wang, G., Tomasi, D., Backus, W., Wang, R., Telang, F., Geliebter, A., Korner, J., Bauman, A., Fowler, J.S., Thanos, P.K., & Volkow, N.D. (2008). Gastric distention activates satiety circuitry in the human brain. *NeuroImage*, *39*, 1824-1831.

Whalen P.J. (2007). The uncertainty of it all. Trends in Cognitive Science, 11, 499-500.

- Wilson-Mendenhall, C.D., Barrett, L.F., Simmons, W.K., & Barsalou, L.W. (2011). Grounding emotion in situated conceptualization. *Neuropsychologia*, 49, 1105-1127.
- Wright, C.E., Strike, P.C., Brydon, L., & Steptoe, A. (2005). Acute inflammation and negative mood: Mediation by cytokine activation. *Brain, Behavior, & Immunity*, 19, 345–350.
- Zaki, J., Davis, J.I., & Ochsner, K.N. (2012). Overlapping activity in anterior insula during interoception and emotional experience. *NeuroImage*, *62*, 493-499.



**Figure 1.** An emergent model of emotion as theorized by the Conceptual Act Theory. Measurable phenomena such as peripheral nervous system activity, visual sensations, episodic knowledge, exogenous attention, etc. contribute to basic components of emotion such as core affect, exteroceptive sensation, conceptual knowledge, and executive function. These components combine in a given context to create an emergent emotion experience. The Conceptual Act Theory proposes that these components combine through the process of *situated conceptualization*.





An emergent model of emotion as theorized by the Conceptual Act Theory. Measurable phenomena such as peripheral nervous system activity, visual sensations, episodic knowledge, exogenous attention, etc. contribute to basic components of emotion such as core affect, exteroceptive sensation, conceptual knowledge, and executive function. These components combine in a given context to create an emergent emotion experience. The Conceptual Act Theory proposes that these components combine through the process of situated conceptualization.

361x270mm (72 x 72 DPI)